

Thermodynamics of Spin-1/2 Heisenberg Chains: Experiment and Theory

D. C. Johnston



Collaborators

Experiment

R. K. Kremer, MPI for Solid State Research, Stuttgart
T. Saito, M. Azuma, M. Takano, Kyoto University
T. Yamauchi, Y. Ueda, University of Tokyo
S. L. Bud'ko, A. F. Panchula, P. C. Canfield, ISU

Theory

M. Troyer, ETH Zürich
X. Wang, IRRNPM, Lausanne
A. Klümper, University of Cologne


Introduction

We have measured the magnetic susceptibility χ and/or specific heat $C_p(T)$ of two insulating compounds containing spin $S = 1/2$ chains: NaV_2O_5 , and the ambient- (AP) and high-pressure (HP) forms of $(\text{VO})_2\text{P}_2\text{O}_7$

NaV_2O_5 is structurally a two-leg spin ladder compound with one electron per V-O-V rung. According to theory, the ladder can be mapped onto a spin-1/2 chain problem. NaV_2O_5 undergoes combined charge, lattice, and spin dimerization transitions at $T_c = 34$ K.

Goals: (1) Measure $\chi(T)$ and $C_p(T)$ on good single crystals of NaV_2O_5 , (2) Compute the temperature-dependent spin gap Δ below T_c from the $\chi(T)$ data and compare the results with those of other measurements such as neutron scattering, (3) From the $C_p(T)$ data, determine the relative contributions of the spin and charge/lattice degrees of freedom to the phase transition at T_c

Structure of NaV_2O_5




$(\text{VO})_2\text{P}_2\text{O}_7$: DCJ discovered in 1987 that this $S = 1/2$ compound has a spin gap, and fitted his $\chi(T)$ data by existing theoretical predictions for $\chi(T)$ of the spin-1/2 AF alternating-exchange chain. The crystal structure suggests that the compound contains two-leg V spin ladders instead, but no predictions for $\chi(T)$ of spin ladders existed at that time 1994: Barnes and Riera fitted DCJ's $\chi(T)$ data by a two-leg ladder model. Inelastic neutron scattering measurements on a powder by Eccleston et al. confirmed the spin ladder model 1997: Garrett et al. ruled out the spin ladder model and re-confirmed the alternating chain model from inelastic neutron scattering data on single crystals. A "troublesome" second spin gap was found, not possible for the alternating chain model, which was attributed to two-magnon bound states 1999: Yamauchi et al. deduced from NMR that $(\text{VO})_2\text{P}_2\text{O}_7$ contains two independent alternating-exchange chains, with the two spin gaps as found in the neutron measurements on single crystals

1999: Azuma et al. discovered a high-pressure phase of $(\text{VO})_2\text{P}_2\text{O}_7$ with a simpler structure containing a single type of alternating-exchange chain

Goals: (1) Measure the $\chi(T)$ of powders and single crystals of the ambient- (AP) and high-pressure (HP) phases of $(\text{VO})_2\text{P}_2\text{O}_7$ and (2) test consistency with the two- and one-chain models, respectively

Structure of $\text{AP}-(\text{VO})_2\text{P}_2\text{O}_7$





Approach

- Carry out numerical calculations of $\chi(T)$ for the $S = 1/2$ uniform and alternating-exchange Heisenberg chain using Bethe ansatz calculations, quantum Monte Carlo (QMC) simulations and transfer-matrix density-matrix renormalization group (TMRG) calculations. Bethe ansatz calculations of the specific heat are also performed for the uniform chain
- Obtain high-precision fits to the numerical data
- Model the experimental data with the theory



Theory

Bethe ansatz $\chi(T)$ and specific heat $C(T)$ data for the AF $S = 1/2$ uniform Heisenberg chain



- The very low-temperature data are in excellent agreement with Lukyanov's 1998 exact field theory predictions

Numerical QMC and TMRG data for AF alternating-exchange chains with various values of $\alpha = J_2/J_1$



- The above set of solid curves is a single high-accuracy two-dimensional fit to the data with $0 \leq \alpha \leq 1$ and $0.01 \leq k_B T/J_1 \leq 5$. This function has been sought by experimentalists for several decades

The $\chi(T)$ function is recast below in terms of the average $J = (J_1 + J_2)/2$ and dimerization parameter $\delta = (J_1 - J_2)/2J$ instead of J_1 and $\alpha = J_2/J_1$. This form is more transparent for modeling $\chi(T)$ data for compounds containing chains with weak alternation and/or that show spin-dimerization transitions

